AETNA BETTER HEALTH® Coverage Policy/Guideline					
Name:	Nuvigil (armodafinil)	Page:	1 of 3	
Effective Date: 6/26/2024			Last Review Date:	6/6/2024	
Applies to:	⊠Illinois □Michigan	□New Jersey □Pennsylvania Kids	□Maryland □Virginia	l	

Intent:

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for Nuvigil (armodafinil) under the patient's prescription drug benefit.

Description:

Nuvigil (armodafinil) is indicated to improve wakefulness in adult patients with excessive sleepiness associated with obstructive sleep apnea (OSA), narcolepsy, or shift work disorder (SWD).

Limitations of Use

In OSA, Nuvigil (armodafinil) is indicated to treat excessive sleepiness and not as treatment for the underlying obstruction. If continuous positive airway pressure (CPAP) is the treatment of choice for a patient, a maximal effort to treat with CPAP for an adequate period of time should be made prior to initiating Nuvigil (armodafinil) for excessive sleepiness.

Applicable Drug List:

Armodafinil

Policy/Guideline:

The requested drug will be covered with prior authorization when the following criteria are met:

• The patient has a diagnosis of narcolepsy

AND

The request is for continuation of therapy

AND

o The patient had a positive response to treatment

OR

- The requested drug is being prescribed by, or in consultation with, a sleep specialist
 AND
- o The diagnosis is confirmed by sleep lab evaluation

OR

The patient has a diagnosis of shift work disorder (SWD)

AND

The request is for continuation of therapy

AND

The patient had a positive response to treatment

AND

The patient is still a shift-worker

	TER HEALTH®	*ae	tna [™]		
Coverage Policy/Guideline					
Name:	Nuvigil (armodafinil)	Page:	2 of 3	
Effective Date: 6/26/2024			Last Review Date:	6/6/2024	
Applies	⊠Illinois	□New Jersey	□Maryland		
to:	□Michigan	□Pennsylvania Kids	□Virginia		

OR

- The requested drug is being prescribed by, or in consultation with, a sleep specialist
- A sleep log and actigraphy monitoring have been completed for at least 14 days and shows a disrupted sleep and wake pattern

AND

• Symptoms have been present for 3 or more months

OR

• The patient has a diagnosis of obstructive sleep apnea (OSA)

AND

The request is for continuation of therapy

AND

The patient had a positive response to treatment

AND

 The patient is compliant with using continuous positive airway pressure (CPAP) or bilevel positive airway pressure (BIPAP)

OR

- The requested drug is being prescribed by, or in consultation with, a sleep specialist
 AND
- The diagnosis has been confirmed by polysomnography

AND

 The patient has been receiving treatment for the underlying airway obstruction (continuous positive airway pressure [CPAP] or bilevel positive airway pressure [BIPAP]) for at least one month

AND

• Treatment with continuous positive airway pressure (CPAP) or bilevel positive airway pressure (BIPAP) will continue

AND

• The patient is unable to take modafinil for the given diagnosis, due to a trial and inadequate treatment response, or intolerance, or a contraindication

Approval Duration and Quantity Restrictions:

Approval: 12 months

Quantity Level Limit:

- Nuvigil (armodafinil) 50 mg: 60 tablets per 30 days
- Nuvigil (armodafinil) 150 mg, 200 mg, 250 mg: 30 tablets / 30 days

References:

1. Nuvigil [package insert]. Parsippany, NJ: Teva Pharmaceuticals; December 2022.

AETNA BETTER HEALTH®						
Coverage Policy/Guideline						
Name:	Nuvigil (armodafinil)		Page:	3 of 3		
Effective Date: 6/26/2024			Last Review Date:	6/6/2024		
Applies to:	⊠Illinois	□New Jersey	□Maryland			
	□Michigan	\square Pennsylvania Kids	□Virginia			

- 2. Lexicomp Online, AHFS DI (Adult and Pediatric) Online. Waltham, MA: UpToDate, Inc.; 2023. https://online.lexi.com. Accessed December 4, 2023.
- 3. Micromedex (electronic version). Merative, Ann Arbor, Michigan, USA. Available at: https://www.micromedexsolutions.com/<u>(cited: 12/04/2023)</u>.
- 4. Kapur VK, Auckley DH, Chowdhuri S, et al. Clinical Practice Guideline for Diagnostic Testing for Adult Obstructive Sleep Apnea: An American Academy of Sleep Medicine Clinical Practice Guideline. *J Clin Sleep Med*. 2017;13(3):479-504.
- 5. Epstein LJ, Kristo D, Strollo PJ, et al. Clinical Guidelines for the Evaluation, Management and Long-term Care of Obstructive Sleep Apnea in Adults. *J Clin Sleep Med.* 2009:5(3):263-276.
- 6. American Academy of Sleep Medicine. International Classification of Sleep Disorders, 3rd edition, text revision. American Academy of Sleep Medicine, 2023.
- 7. Sateia MJ. International Classification of Sleep Disorders- Third Edition: Highlights and Modifications. *CHEST*. 2014;146(5):1387-1394.
- 8. Maski K, Trotti LM, Kotagal S, et al. Treatment of central disorders of hypersomnolence: an American Academy of Sleep Medicine clinical practice guideline. *J Clin Sleep Med.* 2021;17(9):1881-1893.
- 9. Maski K, Trotti LM, Kotagal S, et al. Treatment of central disorders of hypersomnolence: an American Academy of Sleep Medicine systematic review, meta-analysis, and GRADE assessment. *J Clin Sleep Med.* 2021;17(9):1895-1945.