



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Soliris and Biosimilars

Page: 1 of 6

Effective Date: 8/4/2025

Last Review Date: 5/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Michigan
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input checked="" type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input checked="" type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

Intent:

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for Soliris under the patient's prescription drug benefit.

Description:

The indications below including FDA-approved indications and compendial uses are considered covered benefits provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indications

- A. Paroxysmal nocturnal hemoglobinuria (PNH) to reduce hemolysis
- B. Atypical hemolytic uremic syndrome (aHUS) to inhibit complement-mediated thrombotic microangiopathy
- C. Generalized myasthenia gravis (gMG) in adult patients who are anti-acetylcholine receptor (AChR) antibody positive
- D. Neuromyelitis optica spectrum disorder (NMOSD) in adult patients who are anti-aquaporin-4 (AQP4) antibody positive

Bkemv and Epysqli are indicated for the treatment of:

- A. Paroxysmal nocturnal hemoglobinuria (PNH) to reduce hemolysis
- B. Atypical hemolytic uremic syndrome (aHUS) to inhibit complement-mediated thrombotic microangiopathy

Limitations of Use: Soliris, Bkemv, and Epysqli are not indicated for the treatment of patients with Shiga toxin E. coli related hemolytic uremic syndrome (STEC-HUS).

Limitations of Use: Soliris is not indicated for the treatment of patients with Shiga toxin E. coli related hemolytic uremic syndrome (STEC-HUS).

All other indications are considered experimental/investigational and not medically necessary.

Applicable Drug List:

Preferred:
Epysqli

Non-preferred:
Bkemv



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Soliris and Biosimilars

Page: 2 of 6

Effective Date: 8/4/2025

Last Review Date: 5/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Michigan
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input checked="" type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input checked="" type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

Soliris

Policy/Guideline:

Documentation:

Submission of the following information is necessary to initiate the prior authorization review for new requests for treatment of:

A. For initial requests:

1. Atypical hemolytic uremic syndrome: ADAMTS 13 level
2. Paroxysmal nocturnal hemoglobinuria: flow cytometry used to show results of glycosylphosphatidylinositol-anchored proteins (GPI-APs) deficiency
3. Generalized myasthenia gravis: anti-acetylcholine receptor (AChR) antibody positive, clinical classification of myasthenia gravis score, MG activities of daily living score, use of IVIG, use of two immunosuppressive therapies
4. Neuromyelitis optica spectrum disorder: immunoassay used to confirm anti-aquaporin-4 (AQP4) antibody is present

B. For continuation requests: Chart notes or medical record documentation supporting positive clinical response.

Criteria for Initial Approval:

A. Atypical hemolytic uremic syndrome

Authorization of 6 months may be granted for treatment of atypical hemolytic uremic syndrome (aHUS) not caused by Shiga toxin when all of the following criteria are met:

1. ADAMTS 13 activity level above 5%
2. Absence of Shiga toxin
3. The requested medication will not be used in combination with another complement inhibitor (e.g., Ultomiris) for the treatment of aHUS.

B. Paroxysmal nocturnal hemoglobinuria

Authorization of 6 months may be granted for treatment of paroxysmal nocturnal hemoglobinuria (PNH) when all of the following criteria are met:

1. The diagnosis of PNH was confirmed by detecting a deficiency of glycosylphosphatidylinositol-anchored proteins (GPI-APs) (e.g., at least 5% PNH cells, at least 51% of GPI-AP deficient poly-morphonuclear cells)
2. Flow cytometry is used to demonstrate GPI-APs deficiency
3. Member has and exhibits clinical manifestations of disease (e.g., LDH > 1.5 ULN, thrombosis, renal dysfunction, pulmonary hypertension, dysphagia)



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Soliris and Biosimilars

Page: 3 of 6

Effective Date: 8/4/2025

Last Review Date: 5/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Michigan
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input checked="" type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input checked="" type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

4. The requested medication will not be used in combination with another complement inhibitor (e.g., Empaveli, Fabhalta, Piasky, Ultomiris) for the treatment of PNH (concomitant use with Voydeya is allowed).
5. Patient is unable to take the preferred formulary alternative (Epysqli) for the given diagnosis due to a trial and inadequate treatment response or intolerance, or a contraindication.

C. Generalized myasthenia gravis

Authorization of 6 months may be granted for treatment of generalized myasthenia gravis (gMG) when all of the following criteria are met:

1. Anti-acetylcholine receptor (AChR) antibody positive
2. Myasthenia Gravis Foundation of America (MGFA) clinical classification II to IV
3. MG activities of daily living (MG-ADL) total score of greater than or equal to 5
4. Meets one of the following:
 - i. Member has had an inadequate response or intolerable adverse event to at least two immunosuppressive therapies over the course of at least 12 months (e.g., azathioprine, corticosteroids, cyclosporine, methotrexate, mycophenolate, tacrolimus)
 - ii. Member has had an inadequate response or intolerable adverse event to at least one immunosuppressive therapy and intravenous immunoglobulin (IVIG) over the course of at least 12 months
 - iii. Member has a documented clinical reason to avoid therapy with immunosuppressive agents and IVIG
5. The requested medication will not be used in combination with another complement inhibitor (e.g., Ultomiris, Zilbrysq) or neonatal Fc receptor blocker (e.g., Vyvgart, Vyvgart Hytrulo, Rystiggo).

D. Neuromyelitis optica spectrum disorder

Authorization of 6 months may be granted for treatment of neuromyelitis optica spectrum disorder (NMOSD) when all of the following criteria are met:

1. Anti-aquaporin-4 (AQP4) antibody positive
2. Member exhibits one of the following core clinical characteristics of NMOSD:
 - i. Optic neuritis
 - ii. Acute myelitis
 - iii. Area postrema syndrome (episode of otherwise unexplained hiccups or nausea and vomiting)
 - iv. Acute brainstem syndrome
 - v. Symptomatic narcolepsy or acute diencephalic clinical syndrome with NMOSD-typical diencephalic MRI lesions



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Soliris and Biosimilars

Page: 4 of 6

Effective Date: 8/4/2025

Last Review Date: 5/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Michigan
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input checked="" type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input checked="" type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

- vi. Symptomatic cerebral syndrome with NMOSD-typical brain lesions
3. The member will not receive the requested medication concomitantly with other biologics for the treatment of NMOSD.

Continuation of Therapy:

A. Atypical hemolytic uremic syndrome

Authorization of 12 months may be granted for continued treatment in members requesting reauthorization when all of the following criteria are met:

1. There is no evidence of unacceptable toxicity or disease progression while on the current regimen.
2. The member demonstrates a positive response to therapy (e.g., normalization of lactate dehydrogenase (LDH) levels, platelet counts).
3. The requested medication will not be used in combination with another complement inhibitor (e.g., Ultomiris) for the treatment of aHUS.

B. Paroxysmal nocturnal hemoglobinuria

Authorization of 12 months may be granted for continued treatment in members requesting reauthorization when all of the following criteria are met:

1. There is no evidence of unacceptable toxicity or disease progression while on the current regimen.
2. The member demonstrates a positive response to therapy (e.g., improvement in hemoglobin levels, normalization of lactate dehydrogenase [LDH] levels).
3. The requested medication will not be used in combination with another complement inhibitor (e.g., Empaveli, Fabhalta, Piasky, Ultomiris) for the treatment of PNH (concomitant use with Voydeya is allowed).

C. Generalized myasthenia gravis

Authorization of 12 months may be granted for continued treatment in members requesting reauthorization when all of the following criteria are met:

1. There is no evidence of unacceptable toxicity or disease progression while on the current regimen.
2. The member demonstrates a positive response to therapy (e.g., improvement in MG-ADL score, MG Manual Muscle Test (MMT), MG Composite).
3. The requested medication will not be used in combination with another complement inhibitor (e.g., Ultomiris, Zilbrysq) or neonatal Fc receptor blocker (e.g., Vyvgart, Vyvgart Hytrulo, Rystiggo).

D. Neuromyelitis optica spectrum disorder

Authorization of 12 months may be granted for continued treatment in members requesting reauthorization when all of the following criteria are met:



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Soliris and Biosimilars

Page: 5 of 6

Effective Date: 8/4/2025

Last Review Date: 5/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Michigan
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input checked="" type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input checked="" type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

1. There is no evidence of unacceptable toxicity or disease progression while on the current regimen.
2. The member demonstrates a positive response to therapy (e.g., reduction in number of relapses).
3. The member will not receive the requested medication concomitantly with other biologics for the treatment of NMOSD.

Dosage and Administration:

Approvals may be subject to dosing limits in accordance with FDA-approved labeling, accepted compendia, and/or evidence-based practice guidelines.

Approval Duration and Quantity Restrictions:

Approval:

- Initial Requests: 6 months
- Renewals: 12 months

References:

1. Soliris [package insert]. Boston, MA: Alexion Pharmaceuticals, Inc.; March 2024.
2. Loirat C, Fakhouri F, Ariceta G, et al. An international consensus approach to the management of atypical hemolytic uremic syndrome in children. *Pediatr Nephrol*. Published online: April 11, 2015.
3. Parker CJ. Management of paroxysmal nocturnal hemoglobinuria in the era of complement inhibitory therapy. *Hematology*. 2011; 21-29.
4. Sanders D, Wolfe G, Benatar M et al. International consensus guidance for management of myasthenia gravis. *Neurology*. 2021; 96 (3) 114-122 .
5. Jaretzki A, Barohn RJ, Ernstoff RM et al. Myasthenia Gravis: Recommendations for Clinical Research Standards. *Ann Thorac Surg*. 2000;70: 327-34.
6. Hillmen P, Young NS, Schubert J, et al. The complement inhibitor eculizumab in paroxysmal nocturnal hemoglobinuria. *NEJM*. 2006;335:1233-43.
7. Howard JF, Utsugisawa K, Benatar M. Safety and efficacy of eculizumab in anti-acetylcholine receptor antibody-positive refractory generalized myasthenia gravis (REGAIN); a phase 3, randomized, double-blind, placebo-controlled, multicenter study. *Lancet Neurol*. 2017 Oct 20. [http://dx.doi.org/10.1016/S1474-4422\(17\)30369-1](http://dx.doi.org/10.1016/S1474-4422(17)30369-1)Ingenix HCPCS Level II, Expert 2011.
8. Brodsky RA, Young NS, Antonioli E, et al. Multicenter phase 3 study of the complement inhibitor eculizumab for the treatment of patients with paroxysmal nocturnal hemoglobinuria. *Blood*. 2008;111(4):1840-1847.
9. Borowitz MJ, Craig F, DiGiuseppe JA, et al. Guidelines for the Diagnosis and Monitoring of Paroxysmal Nocturnal Hemoglobinuria and Related Disorders by Flow Cytometry. *Cytometry B Clin Cytom*. 2010; 78: 211-230.
10. Preis M, Lowrey CH. Laboratory tests for paroxysmal nocturnal hemoglobinuria (PNH). *Am J Hematol*. 2014;89(3):339-341.
11. Lee JW, Sicre de Fontbrune F, Wong LL, et al. Ravulizumab (ALXN1210) vs eculizumab in adult patients with PNH naive to complement inhibitors: The 301 study. *Blood*. 2018 Dec 3; pii: blood-2018-09-876136.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Soliris and Biosimilars

Page: 6 of 6

Effective Date: 8/4/2025

Last Review Date: 5/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Michigan
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input checked="" type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input checked="" type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

12. Pittock SJ, Berthele A, Kim HJ, et al. Eculizumab in Aquaporin-4-Positive Neuromyelitis Optica Spectrum Disorder. *N Engl J Med*. 2019 May 3. doi: 10.1056/NEJMoA1900866.
13. Wingerchuk DM, Banwell B, Bennett JL, et al. International consensus diagnostic criteria for neuromyelitis optica spectrum disorders. *Neurology*. 2015; 85:177-189.
14. Parker CJ. Update on the diagnosis and management of paroxysmal nocturnal hemoglobinuria. *Hematology Am Soc Hematol Educ Program*. 2016;2016(1):208-216.
15. Dezern AE, Borowitz MJ. ICCS/ESCCA consensus guidelines to detect GPI-deficient cells in paroxysmal nocturnal hemoglobinuria (PNH) and related disorders part 1 - clinical utility. *Cytometry B Clin Cytom*. 2018 Jan;94(1):16-22.
16. Barnett C, Herbelin L, Dimachkie MM, Barohn RJ. Measuring Clinical Treatment Response in Myasthenia Gravis. *Neurol Clin*. 2018 May;36(2):339-353.
17. Bkerv [package insert]. Thousand Oaks, CA: Amgen Inc.; May 2024.
18. Epysqli [package insert]. Republic of Korea: Samsung Bioepis Co., Ltd.; July 2024.